

Limited-time offer! Call Dr. Levens for an appointment today. (954)752-1020

Celebrate and Save

\$50

on a BOTOX[®] Cosmetic treatment

Contact our office today to get your \$50 rebate on the most popular physician-administered aesthetic procedure for 7 years in a row¹

BOTOX[®] Cosmetic
(Botulinum Toxin Type A)
is celebrating:

- 7 years of physician experience
- 11 million procedures and counting²
- High patient satisfaction³

The one. The only.[®]

BOTOX[®]
—Cosmetic
Botulinum Toxin Type A

*Join the 7-Year
Anniversary
Celebration!*

Special \$50 rebate available only through September 30—quantities are limited

Make your appointment today and ask if BOTOX[®] Cosmetic (Botulinum Toxin Type A) is right for you. Don't miss this opportunity to save!

Additional savings also available on future BOTOX[®] Cosmetic treatments. Details provided with your rebate.

*What are you waiting for?
Call Dr. Levens today!*

(954)753-1020

Indication: BOTOX[®] Cosmetic is approved for the temporary treatment of moderate to severe frown lines between the brows in people ages 18 to 65.

Important Safety Information

BOTOX[®] Cosmetic injections should not be given to people who have an infection where the physician proposes to inject. They should not be given to people who are known to be sensitive to any ingredient in BOTOX[®] Cosmetic.

Serious heart problems and serious allergic reactions have been reported rarely. If you think you are having an allergic reaction or other reactions, such as difficulty swallowing, speaking, or breathing, call your doctor immediately. The most common side effects following injection include temporary eyelid droop and nausea. Localized pain, infection, inflammation, tenderness, swelling, redness, and/or bleeding/bruising may be associated with the injection. Patients with certain neuromuscular disorders such as ALS, myasthenia gravis, or Lambert-Eaton syndrome may be at increased risk of serious side effects.

Additional Important Safety Information

The FDA on April 30, 2009, in its update to the early communication sent in 2008, gave the following recommendations:

- Understand that dosage strength (potency) expressed in "Units" or "U" are different among the botulinum toxin products; clinical doses expressed in units are not interchangeable from one botulinum toxin product to another.
- Be alert to and educate patients and caregivers about potential adverse events due to distant spread of botulinum toxin effects following local injections including: unexpected loss of strength or muscle weakness, hoarseness or trouble talking (dysphonia), trouble saying words clearly (dysarthria), loss of bladder control, trouble breathing, trouble swallowing, double vision, blurred vision and drooping eyelids.
- Understand that these adverse events have been reported as early as several hours and as late as several weeks after treatment.
- Advise patients to seek immediate medical attention if they develop any of these symptoms.

Please see accompanying full prescribing information.

1. The American Society of Plastic Surgeons (ASPS). 2009 Report of the 2008 Statistics: National Clearinghouse of Plastic Surgery Statistics. The American Society of Plastic Surgeons (ASPS) Web site. <http://www.plasticsurgery.org/Media/Stats/2008-US-cosmetic-reconstructive-plastic-surgery-minimally-invasive-statistics.pdf>. Accessed June 15, 2009. 2. Data on file, Allergan, Inc.; Safety analysis. 3. Aesthetic Surgery Education and Research Foundation (ASERF). BOTOX[®] Cosmetic Use Survey, April 2005.